

specific antibodies); CD4+, CD8+, and natural killer cells; and functional antibodies to the non-structural protein NS1. We agree that dissection of immune responses over time, including serotype-specific antibodies, in seronegative and seropositive participants in phase 3 efficacy trials is important to understand the role of specific immune responses in protection against each dengue serotype.

Shibadas Biswal, M.D.

Takeda Vaccines
Cambridge, MA
shibadas.biswal@takeda.com

Vianney Tricou, D.Phil.

Takeda Pharmaceuticals International
Zurich, Switzerland

Hansi Dean, Ph.D.

Takeda Vaccines
Cambridge, MA

Drs. Tricou and Dean report being permanent employees of the Takeda Group of companies. Since publication of his article, Dr. Biswal reports no further potential conflict of interest.

1. Dengue haemorrhagic fever: diagnosis, treatment, prevention, and control. 2nd ed. Geneva: World Health Organization, 1997.
2. Rezeki Hadinegoro S, Arredondo-García JL, Capeding MR, et al. Efficacy and long-term safety of a dengue vaccine in regions of endemic disease. *N Engl J Med* 2015;373:1195-206.
3. Kirkpatrick BD, Whitehead SS, Pierce KK, et al. The live attenuated dengue vaccine TV003 elicits complete protection against dengue in a human challenge model. *Sci Transl Med* 2016; 8:330ra36.
4. Tricou V, Low JG, Oh HM, et al. Safety and immunogenicity of a single dose of a tetravalent dengue vaccine with two different serotype-2 potencies in adults in Singapore: a phase 2, double-blind, randomised, controlled trial. *Vaccine* 2020;38:1513-9.
5. Katzelnick, LC, Montoya M, Gresh L, et al. Neutralizing antibody titers against dengue virus correlate with protection from symptomatic infection in a longitudinal cohort. *Proc Natl Acad Sci U S A* 2016;113:728-33.

DOI: 10.1056/NEJMc2000987

Effects of Intermittent Fasting on Health, Aging, and Disease

TO THE EDITOR: In their review article, de Cabo and Mattson (Dec. 26 issue)¹ suggest potential positive effects of intermittent fasting. We would like to highlight safety concerns regarding intermittent fasting in persons with diabetes mellitus. Furmli et al. reported a considerable reduction in or elimination of the need for oral antihyperglycemic agents and insulin in persons with type 2 diabetes after the onset of fasting.² In a prospective trial involving adults with type 2 diabetes who were participating in a 5:2 intermittent fasting regimen (fasting 2 days each week), the mean medication effect score for oral antihyperglycemic agents and insulin was reduced.³ The investigators anticipated a reduction in the dose of the antihyperglycemic agent and used a medication management protocol³ that was modified for patients with hypoglycemia. The dose reduction occurred as early as 5 days after the initiation of the fasting protocol,² and most changes in medications occurred within 3 months.³

When prescribing intermittent fasting for persons with diabetes, a reduction in the dose of the antihyperglycemic agent should be anticipated early in treatment. The risk of hypoglycemia is increased, particularly among patients who are receiving insulin or sulfonylureas, and it is more

pronounced on restricted-feeding days.³ Caution is warranted in persons with a history of hypoglycemic unawareness. Intermittent fasting is probably safe in persons with diabetes that is controlled by dietary changes alone.³

Elizabeth M. Lamos, M.D.

Rana Malek, M.D.

Kashif M. Munir, M.D.

University of Maryland School of Medicine
Baltimore, MD
elamos@som.umaryland.edu

No potential conflict of interest relevant to this letter was reported.

1. de Cabo R, Mattson MP. Effects of intermittent fasting on health, aging, and disease. *N Engl J Med* 2019;381:2541-51.
2. Furmli S, Elmasry R, Ramos M, Fung J. Therapeutic use of intermittent fasting for people with type 2 diabetes as an alternative to insulin. *BMJ Case Rep* 2018;2018:bcr-2017-221854.
3. Carter S, Clifton PM, Keogh JB. Effect of intermittent compared with continuous energy restricted diet on glycemic control in patients with type 2 diabetes: a randomized noninferiority trial. *JAMA Netw Open* 2018;1(3):e180756.

DOI: 10.1056/NEJMc2001176

TO THE EDITOR: The article by de Cabo and Mattson promotes intermittent fasting as a dietary pattern that has cardiovascular benefits, including reduced blood pressure, lipid levels, and inflam-

matory markers. Although the authors also report improvement in insulin sensitivity, the results of one study appear to be variable, especially among patients in different weight categories.¹

Although fasting may affect obese and non-obese patients differently, the particular regimen and how the regimen syncs with one's circadian rhythm may contribute to the success of the diet. Regimens that match the circadian clock may improve insulin sensitivity, lipid production, and blood-pressure control because persons who adhere to such regimens avoid eating late in the day and in doing so may decrease their cardiovascular risk.^{2,3}

Finally, given the benefits that the authors mention with respect to recovery of ischemic tissue after trauma and traumatic brain injury, future studies are warranted to determine whether fasting reduces myocardial injury. A study involving mice with myocardial infarction showed that fasting decreased inflammation, adverse remodeling, and mortality.⁴ Intermittent fasting appears to be promising in reducing cardiovascular risk factors.

Tiffany Dong, M.D.

Melroy D'Souza, M.D.

Laurence Sperling, M.D.

Emory University School of Medicine
Atlanta, GA
lsperli@emory.edu

No potential conflict of interest relevant to this letter was reported.

1. Catenacci VA, Pan Z, Ostendorf D, et al. A randomized pilot study comparing zero-calorie alternate-day fasting to daily caloric restriction in adults with obesity. *Obesity (Silver Spring)* 2016;24:1874-83.

2. Anton SD, Moehl K, Donahoo WT, et al. Flipping the metabolic switch: understanding and applying the health benefits of fasting. *Obesity (Silver Spring)* 2018;26:254-68.

3. Scheer FA, Hilton MF, Mantzoros CS, Shea SA. Adverse metabolic and cardiovascular consequences of circadian misalignment. *Proc Natl Acad Sci U S A* 2009;106:4453-8.

4. Ahmet I, Wan R, Mattson MP, Lakatta EG, Talan M. Cardio-protection by intermittent fasting in rats. *Circulation* 2005;112:3115-21.

DOI: 10.1056/NEJMc2001176

TO THE EDITOR: The review article by de Cabo and Mattson concludes that “intermittent fasting has broad-spectrum benefits for many health conditions,” and the authors present “sample prescriptions” to implement intermittent fasting in clinical practice. However, these clinical con-

clusions and recommendations overstate the evidence presented in the article.

The authors frequently cite results of studies of caloric restriction as evidence of the efficacy of intermittent fasting. Select outcomes are presented from case reports, case series, and small clinical trials to support the idea that “obesity, diabetes, cardiovascular disease, and cancers” are “major indications” for intermittent fasting. The authors present data from three randomized, controlled trials with sample sizes of 100 participants or more that assess intermittent fasting; all three trials compare intermittent fasting with caloric restriction with respect to effects on measures of metabolic health.¹⁻³ Two randomized, controlled trials showed that weight loss and most metabolic markers were not different between participants assigned to intermittent fasting and those assigned to caloric restriction.^{1,2} Although one randomized, controlled trial showed a significantly greater reduction in body fat and insulin resistance with intermittent fasting than with caloric restriction, no significant differences were observed in other metabolic measures, including the glycated hemoglobin level, cholesterol level, weight, and waist circumference.³

Further investigation of intermittent fasting is warranted, given its promising results in animal models. However, it is important that clinical conclusions and recommendations accurately convey the limited clinical evidence available.

Robert A. Kleinman, M.D.

Stanford University
Stanford, CA
rkleinman@stanford.edu

David R. Kleinman, M.D.

University of Alberta
Edmonton, AB, Canada

No potential conflict of interest relevant to this letter was reported.

1. Trepanowski JF, Kroeger CM, Barnosky A, et al. Effect of alternate-day fasting on weight loss, weight maintenance, and cardioprotection among metabolically healthy obese adults: a randomized clinical trial. *JAMA Intern Med* 2017;177:930-8.

2. Harvie MN, Pegington M, Mattson MP, et al. The effects of intermittent or continuous energy restriction on weight loss and metabolic disease risk markers: a randomized trial in young overweight women. *Int J Obes (Lond)* 2011;35:714-27.

3. Harvie M, Wright C, Pegington M, et al. The effect of intermittent energy and carbohydrate restriction v. daily energy restriction on weight loss and metabolic disease risk markers in overweight women. *Br J Nutr* 2013;110:1534-47.

DOI: 10.1056/NEJMc2001176

TO THE EDITOR: In their review article, de Cabo and Mattson describe the eventually favorable assignment of the benefits of caloric restriction to the inadvertent intermittent fasting that participants underwent during caloric restriction. It would be prudent to consider the benefits of intermittent fasting as a possible further surrogate for restriction of protein and especially restriction of branched-chain amino acids (leucine, isoleucine, and valine). These branched-chain amino acids are commonly found in meat products and are difficult to obtain in predominantly plant-based diets.¹

Protein restriction provides a novel route to cellular mechanisms involved in intermittent fasting; in Figure 1 of their article, de Cabo and Mattson show how amino acid sensing can activate the mammalian target of rapamycin (mTOR) directly. In particular, leucine restriction has been shown to have a potent effect on cellular signaling, similar to that which occurs in intermittent fasting.²

Low dietary protein and a predominantly plant-based diet are a fundamental part of the Okinawan diet discussed by the authors. Similar diets are common in so-called blue zones, regions of the world where people have been found to live longer than average.³ Further experimental studies that account for the protein content and, ideally, the amino acid composition of the eating period in intermittent fasting diets are warranted to avoid potentially aberrant conclusions.

Ramy Saad, M.B., B.S.

University of Surrey
Guildford, United Kingdom
ramy.s@hotmail.co.uk

No potential conflict of interest relevant to this letter was reported.

1. Mariotti F, Gardner CD. Dietary protein and amino acids in vegetarian diets: a review. *Nutrients* 2019;11(11):E2661.

2. Dodd KM, Tee AR. Leucine and mTORC1: a complex relationship. *Am J Physiol Endocrinol Metab* 2012;302(11):E1329-E1342.

3. Le Couteur DG, Solon-Biet S, Wahl D, et al. New horizons: dietary protein, ageing and the Okinawan ratio. *Age Ageing* 2016;45:443-7.

DOI: 10.1056/NEJMc2001176

THE AUTHORS REPLY: The letters prompted by our recent review article provide valuable informa-

tion for physicians when they consider the effects of intermittent fasting on the health of patients. Lamos et al. rightly point out that the improvement in insulin sensitivity afforded by intermittent fasting could enable patients with type 2 diabetes to reduce or eliminate their dependence on antihyperglycemic drugs. We agree that it is imperative that these patients transition to an intermittent-fasting eating pattern under the supervision of their physicians and with careful monitoring of blood glucose levels while tapering their medication dose. The potential for hypoglycemia during short fasting periods can be reduced by a gradual transition to intermittent fasting, as suggested in Figure 4 of our article.

In the case of daily time-restricted eating, the question posed by Dong et al. of whether and how the timing window for eating and circadian rhythms affect health outcomes remains to be answered. This is important because the notion that breakfast is the “most important meal” is not supported by evidence.¹ To properly answer this question, randomized, controlled trials that are designed to directly compare early with late 6-to-8-hour eating periods should last at least 2 months to enable full adaptation to the new eating pattern.

In response to Kleinman and Kleinman: our review article stated that animals on caloric restriction consume their entire daily allotment of food within a few hours after its provision and, therefore, they fast for approximately 20 hours every day (see Section S1 in the Supplementary Appendix, available with the full text of our article at NEJM.org).^{2,3} Regarding studies in humans, we agree that randomized, controlled trials of intermittent fasting should include a control group matched for weekly calorie intake.⁴ Our suggestion that intermittent fasting might prove to be beneficial in patients with diabetes, cardiovascular disease, and cancer is based on clear findings from studies in animals and from scores of studies in humans.

Finally, Saad notes that inhibition of the mTOR pathway by dietary protein restriction may elicit beneficial effects on health similar to those of intermittent fasting. This has important implications for health and disease risk because the notion that a high-protein intake is good for health is a myth promulgated by some food and dietary-supplement industries. However, a low-

protein diet alone may not elicit the same organismal response as intermittent fasting, and data from randomized, controlled trials that compare intermittent fasting with low-protein diets are lacking.

Mark P. Mattson, Ph.D.

Johns Hopkins University School of Medicine
Baltimore, MD
mrmattso2@jhmi.edu

Rafael de Cabo, Ph.D.

National Institute on Aging
Baltimore, MD

Since publication of their article, the authors report no further potential conflict of interest.

1. Casazza K, Fontaine KR, Astrup A, et al. Myths, presumptions, and facts about obesity. *N Engl J Med* 2013;368:446-54.
2. Acosta-Rodriguez VA, de Groot MHM, Rijo-Ferreira F, Green CB, Takahashi JS. Mice under caloric restriction self-impose a temporal restriction of food intake as revealed by an automated feeder system. *Cell Metab* 2017;26(1):267-277.e2.
3. Mitchell SJ, Bernier M, Mattison JA, et al. Daily fasting improves health and survival in male mice independent of diet composition and calories. *Cell Metab* 2019;29(1):221-228.e3.
4. Harvie M, Wright C, Pegington M, et al. The effect of intermittent energy and carbohydrate restriction v. daily energy restriction on weight loss and metabolic disease risk markers in overweight women. *Br J Nutr* 2013;110:1534-47.

DOI: 10.1056/NEJMc2001176

Correspondence Copyright © 2020 Massachusetts Medical Society.

INSTRUCTIONS FOR LETTERS TO THE EDITOR

Letters to the Editor are considered for publication, subject to editing and abridgment, provided they do not contain material that has been submitted or published elsewhere.

Letters accepted for publication will appear in print, on our website at NEJM.org, or both.

Please note the following:

- Letters in reference to a *Journal* article must not exceed 175 words (excluding references) and must be received within 3 weeks after publication of the article.
- Letters not related to a *Journal* article must not exceed 400 words.
- A letter can have no more than five references and one figure or table.
- A letter can be signed by no more than three authors.
- Financial associations or other possible conflicts of interest must be disclosed. Disclosures will be published with the letters. (For authors of *Journal* articles who are responding to letters, we will only publish new relevant relationships that have developed since publication of the article.)
- Include your full mailing address, telephone number, fax number, and email address with your letter.
- All letters must be submitted through our online submission system at NEJM.org.

Letters that do not adhere to these instructions will not be considered. We will notify you when we have made a decision about possible publication. Letters regarding a recent *Journal* article may be shared with the authors of that article. We are unable to provide prepublication proofs. Submission of a letter constitutes permission for the Massachusetts Medical Society, its licensees, and its assignees to use it in the *Journal's* various print and electronic publications and in collections, revisions, and any other form or medium.

THE JOURNAL'S WEB AND EMAIL ADDRESSES

To submit a letter to the Editor: authors.NEJM.org

For information about the status of a submitted manuscript: authors.NEJM.org

To submit a meeting notice: meetingnotices@NEJM.org

The *Journal's* web pages: NEJM.org